

Genetic variation among cat fish (*Mystus carcio*) population assessed by randomly amplified polymorphic (RAPD) markers from Assam, India

Innifa Hasan¹ and Mrigendra Mohan Goswami²

1. Department of Zoology, Handique Girls' College, Panbazar, Guwahati-781001

2. Department of Zoology, Gauhati University, Guwahati-781007

Corresponding author: Innifa Hasan, Department of Zoology, Handique Girls' College,

Reviewer: 1) Dr Diganta goswami, Defence Research Laboratory, Scientist, Tezpur Assam

2) Dr Hiranmoyee Sharma, Assistant Professor, D.N College, Arunachal Pradesh

Abstract

Catfishes are among the most studied group of vertebrates in aquaculture today as they provide a good model system for studying the evolution of vertebrate behavior. In the present study, random amplified polymorphic DNA polymerase chain reaction (RAPD-PCR) analysis was applied to find the phylogenetic relationship among the individuals of *Mystus carcio* populations. The present finding revealed that different primers have different performances for evaluation of genetic polymorphism. Total 412 RAPD fragments were generated using nine decamer primers of arbitrary nucleotide sequences. In the experiment 322 polymorphic bands and 90 monomorphic bands were produced which shows 78.15 % of polymorphism and 21.84 % of monomorphism. UPGMA dendrogram constructed on the basis of genetic distance formed three distinct clusters indicating comparatively higher level of genetic variations in the studied *M. carcio* populations in Assam. Once the population structure is known, scientific management for optimal harvest and conservation of the catfish fishery resource can be undertaken. Therefore, the present study may serve as a reference for future examinations of genetic variations within the populations of fishes which are commercially important and the possible use of DNA markers in future may create new avenues for cat fish molecular biological research in this part of world.

Key words: Genetic, diversity, RAPD, markers, cat fish.

Introduction:

Catfishes are among the most studied group of vertebrates in aquaculture today as they provide a good model system for studying the evolution of vertebrate behavior. *Mystus carcio* is distinguished from its congeners in having a shorter adipose fin base, a long posterior fontanel extending to the base of occipital process, anterior fontanel extending from the level of posterior margin of nasal opening through posterior orbital margin; supra occipital process narrow and extending upto basal bone of dorsal fin. This important species for aquaculture is naturally distributed in India, West Bengal and Assam (Ganga-Brahmaputra drainages), Bangladesh: Meghana basin in Comilla district. (Darshan et al., 2011). Even though the catfishes are in great demand in the Indian domestic markets, the catfish aquaculture including *Mystus* sp has not yet been developed for its aquaculture potential (Tripathi 1996). The entire demand for this fish in the domestic market is met through capture from river bodies and hence the effective management of wild stocks is critical. For the development of effective management strategies information on population structure is essential to conserve the biodiversity associated with different species, sub-species, stocks and races (Turan et al., 2005). Therefore for good aquacultural practices and to maintain a healthy gene pool, detailed knowledge on the population structure of *Mystus* sp. is needed.

All organisms are subject to mutations because of normal cellular operations or interactions with the environment, leading to genetic variation (polymorphism). Genetic variation in a species enhances the capability of organism to adapt to changing environment and is necessary for survival of the species (Fisher, 1930). Genetic variation arises between individuals leading to differentiation at the level of population, species and higher order taxonomic groups apart from other evolutionary forces like selection and genetic drift. Molecular markers along with the development of new statistical tools has revolutionized the analytical power necessary to explore the genetic diversity, both in native populations and in captive lots (Tamanna et al., 2012). Nowadays, a wide range of new molecular techniques have been explored and reported for fishes (Lehmann et al., 2000; Jayasankar, 2004). Random amplified polymorphic DNA (RAPD) one such technique which was first introduced by Williams et al. (1990). RAPD technique is the one of the most frequently used molecular methods for taxonomic and systematic analyses of various organisms and has provided important applications in catfish (Bartish et al., 2000). RAPD has also been used to estimate genetic diversity and variations needed to study fish management and conservation practices, even with endangered species (Lopera-Barrero et al., 2006 ; Shair et al., 2011).

It is based on the PCR amplification of discrete regions of genome with short oligonucleotide primers of arbitrary sequence (Welsh and McClelland, 1990; Williams et al., 1990). The characters assessed through RAPD are useful for genetic studies because they provide various types of data-taxonomic population, inheritance pattern of various organisms including fishes (David and Pandian, 2006).

The information on morphometric measurements of fishes and the study of statistical relationship among them are essential for taxonomic work (McConnel, 1978). Moreover, to know the origin of stock, separation of stocks or identification of

commercially important species of fishes, morphometric characters are frequently used (Royce, 1963; Kramholz and Cavanah, 1968). These type of study are important for understanding the interactive effect of environment, selection and heredity on the body shapes and sizes within a species (Cadrin 2000). Several studies on the comparative morphometrics of different fish populations have been conducted (Ibanez- Aguirre *et al.*, 2006). The study of genetic diversity of catfishes of Assam is very much limited, so in the present study, this technique was applied to analyze the genetic relationship among *Mystus carcio* populations. The objectives of this study are focused on morphometric identification and detection of RAPD pattern for determination of the genetic variation of a population of *Mystus carcio* from Assam, India.

Materials and methods:

Fish sampling sites and morphometric measurements of fishes:

Geographically, populations of *Mystus carcio* were caught from freshwater bodies of Assam about 100-400 km away from each other, that is, Uzan Bazar (Brahmaputra river, Kamrup District) ; Kolong river at Morigaon (Morigaon District) ; Dhemaji local fish market (Dhemaji District) in the month of June, 2013. A total of 30 fish specimens were collected from all the locations with the help of local fishermen and 20 fish specimens were randomly selected for morphometric measurements and estimating genetic variations. All the fish specimens were kept in the iceboxes and brought to the laboratory for further study. For the morphometric measurements, total 27 parameters were considered. Fish specimens were morphologically identified with taxonomic keys (Srivastava, 2000; Jayaram, 2002). The muscle tissues were isolated from freshly caught fishes and preserved at -20°C for further use.

Isolation of Genomic DNA from Fish Tissue:

For the isolation of total genomic DNA, a modified protocol was followed using Sambrook and Russel(Molecular Cloning A Lab. Manual). UV VIS spectrophotometer was used to check quality as well as quantity of isolated DNA. Optical densities of the DNA samples were measured at 260 nm and 280nm and the concentration of extracted DNA was adjusted to 50ng/ µl for PCR amplification.

PCR primers:

In the present study, 30 commercially available RAPD primers (10 to 20 base long) made by Xcelris Genomics, India were used to initiate PCR amplifications. Primers were randomly selected on the basis of GC content and annealing temperature for RAPD-PCR amplifications.

PCR amplification:

The reaction mixture (10 µl) for PCR was composed of 1 µl of 10 X Taq polymerase buffer, 1 µl of 2.5 mM dNTPs, 1 µl of RAPD primer, 0.15 µl Taq DNA polymerase (2 U/ µl), 5.55 µl PCR grade water, 0.3 µl of 50mM MgCl₂ and 1µl template DNA. A negative control, without template DNA was also included in each round of reactions.

After preheating for 5 mins at 94°C, PCR was run for 35 cycles. It consisted of a 94°C denaturation step (1min), 37°C annealing step (1 min) and 72°C elongation step (2 min) in a thermal cycler (Biorad). At the end of the run, a final extension period was appended (72°C, 10 min) and then stored at 4°C until the PCR products were analyzed.

Agarose gel electrophoresis:

The amplified DNA fragments were separated on 1.8% agarose gel and stained with Ethidium bromide. A low range DNA marker of 100 bp from Bangalore Genei, Bangalore, India was run with each gel. The amplified pattern was visualized on an UV transilluminator and photographed by gel documentation system (BIORAD, USA).

Statistical analysis:

The RAPD fragments were scored for the presence and absence of fragments on the gel photographs and RAPD fragments were compared among the *M. carcio* populations. RAPD banding patterns were recorded on spreadsheets, which were used to determine gene diversity, gene flow, number of polymorphic loci and genetic distance through a construct by an un-weighted pair group method of arithmetic mean or UPGMA (Nei, 1978) using GGT 2.0 software (<http://www.dpw.wau.nl/pv/pub/ggt/>)

The similarity index (SI) values between the RAPD profiles of any 2 individuals on the same gel were calculated using following formula:

$$\text{Similarity Index (SI)} = 2 N_{AB} / (N_A + N_B)$$

Where,

N_{AB} = total number of RAPD bands shared in common between individuals A and B

N_A = total number of bands scored for individual A

N_B = total number of bands scored for individual B (Lynch, 1990).

Cluster analysis was carried out using GGT 2.0 version software. Dendograms were constructed by employing UPGMA (Unweighted Pair Group Method with Arithmetic Mean) based on Sneath and Sokal(1973) to study the genetic variability within the species. Similarly the same method was followed to construct the dendogram to study the phylogenetic relationship among the genotypes of *Mystus carcio*.

Results and Discussion:

The morphometric variation among the different individuals of *Mystus carcio* was found to be very low (Table-1).

Table1: Morphometric measurements of *Mystus carcio*.

%SL	Range(mm)	Mean \pm SD
Body depth	19.3-30.5	23.4 \pm 6.1
Head length	24.7-30.5	27.3 \pm 2.9
caudal peduncle length	13.9-16.7	15.2 \pm 1.4
Caudal peduncle depth	10.5-13.5	12.2 \pm 1.5
Dorsal to Adipose Distance	10.5-12.4	11.4 \pm 0.9
Dorsal fin base length	14.7-16.0	15.3 \pm 0.6
Dorsal fin length	16.6-18.1	17.2 \pm 0.8
Adipose fin base length	10.6-13.0	11.9 \pm 1.2
Pectoral fin length	18.0-22.0	20.2 \pm 2.0
Pelvic fin length	13.1-16.2	14.8 \pm 1.6
Anal fin length	19.4-21.0	20.1 \pm 0.8
Caudal fin length	26.0-29.0	27.4 \pm 1.5
Pre dorsal length	38.6-48.0	43.2 \pm 4.7
Pre pelvic length	49.0-52.9	50.6 \pm 2.0
Pre anal length	68.1-74.7	70.9 \pm 3.4
Post adipose length	16.6-19.0	17.9 \pm 1.2
%HL		
Head depth	69.2-72.3	70.9 \pm 1.6
Head width	74.0-76.9	75.3 \pm 1.5
Snout length	30.5-35.3	32.9 \pm 2.4
Eye diameter	20.4-23.3	21.9 \pm 1.5
Inter orbital distance	36.0-40.3	38.4 \pm 2.2
Wide of gap of mouth	37.0-42.0	39.4 \pm 2.5
Inter nostril distance	10.8-12	11.3 \pm 0.6
Nasal barbel length	51.7-58.1	54.9 \pm 3.2
Maxillary barbel length	140.6-160.4	152.1 \pm 10.3
Inner mandibular barbel Length	47.6-58.3	52.0 \pm 5.6
Outer mand. Barbel L	95.7-98.4	97.1 \pm 1.4

RAPD polymorphisms:

Among the 30 primers initially tested, only nine R-4, R-5, R-6, R-11, R-12, R-13, R-20, R-21 and R-22 were selected that yielded relatively large number of good quality bands. All the primers produced different RAPD patterns, and the number of fragments amplified per primer varied. The nine primers yielded a total of 412 reproducible and consistently scorable RAPD bands of which 322 were found to be polymorphic and 90 were monomorphic. The number of bands per primer ranged from 5 to 13. Among the primers, R-4 and R-21 gave DNA profile with highest number of bands while R-13 gave the least (Table 3). The RAPD profile of the bands

obtained in the population of *M. carcio* with primer R-4,R-5,R-6 and R-11 is shown in the Figure1 as representative photographs. The UPGMA dendrogram was prepared based on genetic distance by the GGT 2.0 software.The unweighted dendrogram divided all the genotypes in three clusters.




Fig1: RAPD profile of *Mystus carcio* with primers R4, R5, R6 and R11




Fig 2: Genetic polymorphisms in 10 genotypes of *Mystus carcio*




Fig 3: Genetic similarity in 10 genotypes of *Mystus carcio*

Table 2: Pattern of polymorphism (primer wise) in 10 individuals of *Mystus carcio*

Sl no	Polymorphism	<i>Mystus carcio</i>								
		R-4	R-5	R-6	R-11	R-12	R-13	R20	R21	R22
1	Total no of bands	58	36	38	48	37	35	53	58	49
2	Total no of polymorphic bands	48	26	28	38	27	25	43	48	39
3	Total no of monomorphic bands	10	10	10	10	10	10	10	10	10
5	Polymorphism(%)	82.75	78.8	73.6	79.16	72.9	71.4	81.1	82.7	79.5
6	Monomorphism(%)	18.18	27.7	26.3	27.7	27.1	28.5	18.8	17.2	20.5

Little morphological differences were revealed among the *M. carcio* populations by morphometric and meristic studies that were sampled from different rivers and localities in Assam. Most of the morphometric characteristics of the fishes in the present study were similar and there is overlap in the range of each of traditional morphometric measurements taken (Table 1) The morphometric data may not be enough to support the established genetic structure of the population that often leads to taxonomic uncertainty in many occasions because of the considerable geographical and ecological variability in form (Ponniah and Gopalkrishnan,2000; Garg et al.,2009b). In *Hilsa* sp., significant differences in allele frequencies and morphological variations were observed from nine different sites within Bangladesh

as reported by Salini et al., 2004 which may be due to the local environmental conditions.

DNA fingerprinting method has tremendous potential in aquaculture and in fisheries as a tool for identification of individuals (Jong-Man, 2001) and population genetics studies. (Bielawski and Pumo, 1997; Smith et al., 1997; Mamuris et al., 1998). RAPD-PCR is a useful tool for estimating the genetic variability and degree of similarity among fish species as has been reported by other workers (Barman et al., 2003; Jayasankar et al., 2004; Lopera-Barrero et al., 2006; Shair et al., 2011). Using a RAPD analysis, the intrapopulation variation was detected with different primers in tilapia (Bardakci and Skibinski, 1994). Chong et al. in 2000 could identify and characterize *Mystus nemurus* populations by RAPD analysis in Malaysia. RAPD technique has also been used to determine genetic variation within and among three populations of *Mystus vittatus* by Tamanna et al. in Bangladesh in the year 2012.

In the present study among the 30 single decamer random primers, nine primers generated a total of 412 bands in the population which were found to be both polymorphic and monomorphic. In the experiment 322 polymorphic bands and 90 monomorphic bands were produced which shows 78.15 % of polymorphism and 21.84 % of monomorphism (Table-2). Polymorphism for genetic similarity among the different individuals of *M. vittatus* which was analyzed using GGT software is expressed in Fig-2. The cluster analysis and dendrogram showing genetic relationship between 10 genotypes of *M. vittatus* showed formation of 3 clusters (Fig-3). Cluster I include genotype 4, 7, 9 and 8; Cluster II include genotype 10, 3 and 1; Cluster III include genotype 6, 5 and 2.

The present study indicates that comparatively higher level of genetic variation exists in the studied *M. carcio* populations in Assam. Garg et al., 2009a could discriminate between the different populations of *M. vittatus* in reservoirs of Madhya Pradesh by RAPD analysis. According to them the intraspecific genetic similarity between the individuals of the population was due to geological variations or changes in aquatic environment. Our statistical analysis showed considerable genetic variation among the genotypes of *M. carcio* collected from different locations of Assam. This population genetic differentiation may be due to ecological, geographical and evolutionary factors. The genetic diversity data has varied applications in research on evolution, conservation and management of natural resources and genetic improvement programmes, etc. RAPD analysis for genetic diversity study provides a basis to obtain genetic variation within and among populations. Once the population structure is known, scientific management for optimal harvest and conservation of the catfish fishery resource can be undertaken. Therefore, the present study may help researchers serve as a reference point for future examinations of genetic variations within the populations of fishes which are commercially important and the possible use of DNA markers in future may create new avenues for fish molecular biological research.

Acknowledgement

Authors are thankful to the Institutional Biotech Hub, Handique Girls' College and

UGC-SAP(DRF) phaseII Aquaculture Project, Zoology Department, Gauhati University for providing laboratory facilities with all molecular biological tools for executing this research work.

REFERENCES:

1. Barman et al., 2003. Genetic variation between four species of Indian carp as revealed by random amplified polymorphic DNA assay. *Aquaculture.*, 217: 115-123.
2. Bartish IV, Gorkava LP, Rumpunen K, Nybom H 2000. Phylogenetic relationship and differentiation among and within populations of *Chaenomeles* Lindl. (Rosaceae) estimated with RAPD's and isozymes. *Theor. Appl. Genet.* 101: 554-563.
3. Bielawski JP, Pumo DE 1997. Randomly Amplified Polymorphic DNA (RAPD) Analysis of Atlantic coast Striped bass. *Heredity*, 78: 32-40.
4. Cadrin SX 2000. Advances in morphometric identification of fishery stocks. *Reviews in Fish Biology and Fisheries* 10, 91–112
5. Chong,L.K, Tan,S.G, Yusoff,K and Siraj,S.S. 2000. Identification and Characterization of Malaysian River Catfish, *Mystus nemurus* (C&V): RAPD and AFLP Analysis. *Biochemical Genetics, Vol. 38, Nos. 3/4, 2000*
6. Darshan A, Anganthoibi N and Vishwanath W 2011. Redescription of the striped catfish *Mystus carcio* (Hamilton) (Siluriformes: Bagridae), *Zootaxa* 2475: 48-54
7. David, C.J. and Pandian, T.J. 2006. GFP reporter gene confirms paternity in the androgenote Buenos Aires tetra, *Hemigrammus muscaudovittatus*. *J. Exp. Zool.* 305A: 83-95.
8. Fisher, R.A. 1930. *The Genetical Theory of Natural Selection*. Oxford University Press, UK.
9. Garg, R.K., Silawat, N., Sairkar, P., Vijay, N. and Mehrotra, N.N 2009a. RAPD analysis for genetic diversity of two populations of *Mystus vittatus* (Bloch) of Madhya Pradesh, India. *Afric. J. Biotechnol.*, 8 (17): 4032-4038.
10. Garg, R.K., Silawat, N., Sairkar, P., Vijay, N. and Mehrotra, N.N. 2009b. Genetic diversity between two populations of *Heteropneustes fossilis* (Bloch) using RAPD profile. *Int. J. Zool. Res.*, 5: 171-177.
11. Ibanez-Aguirre AL, Cabral-Solis E, Gallardo-Cabello M, Espino-Barr E 2006. Comparative morphometrics of two populations of *Mugil curema* (Pisces: Mugilidae) on the Atlantic and Mexican Pacific coasts. *Scientia Marina* 70 (1), 139–45.
12. Jayaram K.C 2002. *The Fresh Water Fishes of the Indian Region*. Narendra Publishing House, Delhi.
13. Jayasankar P 2004. Random amplified polymorphic DNA (RAPD) fingerprinting resolves species ambiguity of domesticated clown fish (genus: *Amphiprion*, family: Pomacentridae) from India. *Aquacult. Res.*, 35: 1006-1009

14. Jong-Man Y 2001. Genetic similarity and difference between common carp and Israeli carp (*Cyprinus carpio*) based on Random Amplified Polymorphic DNA analyses. *Kor. J. Biol. Sci.* 5: 333-339.
15. Kramholz, C. and Cavanah, F. 1968. Comparative morphometry of fresh water drum from two mid-western localities. *Trans. Am. Fish. Soc.*, **97**: 429-441.
16. Lehmann, J., Stohr, T. and Feldon, J. 2000. Long-term effects of prenatal stress experiences and postnatal maternal separation on emotionality and attentional processes. *Beh. Brain Res.* 107: 133-144.
17. Lopera-Barrero et al., 2006. Genetic diversity in piracanjuba populations (*Brycon orbignyanus*) with the RAPD (Random Amplified Polymorphic DNA) markers. *Journal Animal Science.*, 84:170.
18. Lynch, M. 1990. The similarity index and DNA fingerprinting. *Mol. Biol. Evol.*, 7: 478-484.
19. Mamuris Z, Apostolitis AP, Theodorou AJ, Triantaphyllidis C 1998. Application of random amplified polymorphic DNA (RAPD) markers to evaluate intraspecific genetic variation in red mullet (*Mullus barbatus*). *Mar. Biol.* 132: 171-178
20. McCONNEL, R.H.L., 1978. Identification of fresh water fishes. In: *Methods for assessment of fish production in fresh water* (eds. T. Begenal), pp. 46-83. Blackwell Scientific Publications, London
21. Nei M 1978. Estimation of average heterozygosity and genetic distances from small number of individuals. *Genetics*, 89: 583-590.
22. Ponniah AG, Gopalakrishan A 2000. Cultivable, ornamental, sport and food fishes endemic to Peninsular India with special references to Western Ghats. In: Ponniah AG and Gopalakrishan A (eds.). *Endemic fish diversity of Western Ghats*. NBFGR-NATP Publication, National Bureau of Fish Genetic Resources, Lucknow, India. pp. 13-32.
23. Royce, W.F., 1963. A morphometric study of yellow fin tuna *Thunnus albacore* (Bonnatere) U.S. Fish & Wildlife Services, *Fishery Bull.*, **63**: 395-443.
24. Sambrook, J. and Russell, D.W. 1989. *Molecular cloning: A laboratory manual*, 2nd edition. 1-3 Cold Spring Harbor Laboratory Press, New York.
25. Shair OHM, Al- Ssum R.M, and Bahkali A.H 2011. Genetic variation investigation of tilapia grown under Saudi Arabian controlled environment. *Am. J. Biochem. Mol. Biol.*, 1: 89-94
26. Sneath P. H. A. and Sokal R. R. 1973. *Numerical taxonomy*. Freeman, San Francisco.
27. Smith P.J, Benson P.G, Mcveagh S.M 1997. Comparison of three genetic methods used for stock discrimination of orange *Hoplostethys atlanticus*: Alloenzyme mitochondrial DNA and Random Amplified Polymorphic DNA. *Fish. Bull.* 95: 800-811.
28. Salini J.P, Milton DA, Rahman M.J, Hussain M.G 2004. Allozyme and morphological variation throughout the geographical range of the tropical Hilsa shad *Tenualosa ilisha*. *Fish. Res.*, 55: 53-69.
29. Srivastava C.B.L. 2000: *A Text book of fishery science and Indian Fisheries Kitab Mahal*; 28, Netaji Subhas Road, New Delhi-110 002.

30. Tamanna F.M, Rashid J, Alam M.S. 2012. High levels of genetic variation revealed in wild and hatchery populations of the stripped dwarf catfish *Mystus vittatus* (Bloch) (Bagridae:Siluriformes) in Bangladesh by Random Amplified Polymorphic DNA techniques. *Int J Adv Biol Res* 2(2), 322-327.
31. Tripathi S.D 1996. Present status of breeding and culture of catfishes in South Asia. *Aquatic Living Resources* 9, 219–28.
32. Turan C, Yalcin S, Turan F, Okur E, Akyurt I 2005. Morphometric comparisons of African catfish, *Clarias gariepinus*, populations in Turkey. *Folia Zoologica* 54 (1-2), 165–72.
33. Talwar P.K, Jhingran A.G 1991. Inland fisheries of India and adjacent countries, vol 2. Oxford and IBH Publishing Co, New Delhi, India, pp 545–575
34. Welsh, J. and Mc Clelland, M. 1990. Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Res.*, 18: 7213-7218.
35. Williams, J.G.K., Kubelik, A.R., Livak, K.J., Rafalski, J.A. and Tingey, S.V. 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res.* 18: 6531-6535.

